

# *Drosophila* molting neurohormone bursicon is a heterodimer and the natural agonist of the orphan receptor DLGR2<sup>☆</sup>

Fernando M. Mendive<sup>a,1</sup>, Tom Van Loy<sup>b,1</sup>, Sylvie Claeysen<sup>a,2</sup>, Jeroen Poels<sup>b</sup>, Michael Williamson<sup>c</sup>, Frank Hauser<sup>c</sup>, Cornelis J.P. Grimmelikhuijzen<sup>c</sup>, Gilbert Vassart<sup>a,d,\*</sup>, Jozef Vanden Broeck<sup>b,\*</sup>

<sup>a</sup> *Institut de Recherche Interdisciplinaire en Biologie Humaine et Moléculaire (IRIBHM), Université Libre de Bruxelles, Campus Erasme, 808 Route de Lennik, B-1070 Bruxelles, Belgium*

<sup>b</sup> *Laboratory for Developmental Physiology, Genomics and Proteomics, Zoological Institute, Naamsestraat 59, B-3000 Leuven, Belgium*

<sup>c</sup> *Department of Cell Biology and Comparative Zoology, Institute of Biology, University of Copenhagen, Universitetsparken 15, DK-2100, Copenhagen, Denmark*

<sup>d</sup> *Department of Medical Genetics, Université Libre de Bruxelles, Hôpital Erasme, 808 Route de Lennik, B-1070 Bruxelles, Belgium*

Received 23 February 2005; revised 2 March 2005; accepted 2 March 2005

Available online 17 March 2005

Edited by Robert Barouki

**Abstract** Bursicon is a neurohumoral agent responsible for tanning and hardening of the cuticle and expansion of the wings during the final phase of insect metamorphosis. Although the hormonal activity was described more than 40 years ago, the molecular nature of bursicon has remained elusive. We identify here *Drosophila* bioactive bursicon as a heterodimer made of two cystine knot polypeptides. This conclusion was reached in part from the unexpected observation that in the genome of the honey bee, the orthologs of the two *Drosophila* proteins are predicted to be fused in a single open reading frame. The heterodimeric *Drosophila* protein displays bursicon bioactivity in freshly eclosed neck-ligated flies and is the natural agonist of the orphan G protein-coupled receptor DLGR2.

© 2005 Federation of European Biochemical Societies. Published by Elsevier B.V. All rights reserved.

**Keywords:** Bursicon; G protein-coupled receptor; Leucine-rich repeats containing GPCR; Cystine knot; Comparative genomics; Glycoprotein hormone receptor

## 1. Introduction

Bursicon was described more than 40 years ago as a bioactivity present in CNS (central nervous system) extracts and hemolymph of insects after eclosion, capable of rescuing flies from defective tanning and sclerotization of the cuticle caused by neck-ligation [1–4]. It is a key regulator of the final steps in the stereotyped developmental and behavioural program of ecdysis, i.e., the shedding of the remains of the old cuticle and the consequent maturation of the new one. Bursicon is believed to act downstream of a sequence of regulatory peptides, such as ecdysis triggering hormone (ETH), eclosion hormone (EH) and crustacean cardioactive peptide (CCAP) [5–11]. Bursicon bioactivity can be partially mimicked by cyclic AMP analogs [12,13] and was recently shown to be involved in apoptotic removal of wing epidermal cells during the last step of metamorphosis into adult *Drosophila* [14]. Purification of bioactive bursicon from nerve cords of the cockroach *Periplaneta americana* yielded a 30 kDa protein [15] from which several tryptic peptides were obtained and micro-sequenced [16]. Comparison of these sequences with available genomic data identified a 522 nucleotide open reading frame (ORF) in *Drosophila* (CG13419) that codes for a cystine knot protein with close orthologs in the bee *Apis mellifera* and in the mosquito *Anopheles gambiae* [17,18] (Fig. 1A). The link between the *Drosophila* ORF and bursicon was established from the defective post-ecdysial behaviour of flies with loss-of-function mutations in the putative gene, and the corresponding decrease in bursicon bioactivity in mutant extracts [17]. Accordingly, it was proposed that bursicon would be a homodimeric 30 kDa protein made of two 15 kDa CG13419 mature protomers [17].

DLGR2 is a G protein-coupled receptor of *Drosophila* distantly related to the vertebrate glycoprotein hormone and relaxin receptors [19,20]. It has been proposed as the receptor for bursicon because flies with loss-of-function mutations in DLGR2 display a phenotype (*rickets rk<sup>-/-</sup>*) which is similar to bursicon deficiency induced by neck-ligation. It consists in defective sclerotization and melanization of the cuticle, as well as abnormal wing expansion after adult eclosion. These mutants produce bioactive bursicon, and their phenotype cannot be rescued by bursicon injection [13].

<sup>☆</sup> This study was initially submitted to a major science journal on December 8th 2004, and finally rejected on February 16th 2005 on the basis that a competing, confirmatory study by Luo et al. was published on-line in PNAS on February 10th (submitted 17 December, accepted 31 December) [31].

\*Corresponding authors. Fax: +32 2 555 4212.  
E-mail addresses: gvassart@ulb.ac.be (G. Vassart),  
Jozef.VandenBroeck@bio.kuleuven.ac.be (J. Vanden Broeck).

<sup>1</sup> These authors contributed equally.

<sup>2</sup> Present address: Centre CNRS-INSERM de Pharmacologie-Endocrinologie, CNRS, UPR 9023, 141 rue de la Cardonille, 34094 Montpellier cedex 5, France.

**Abbreviations:** ORF, open reading frame; LGR, leucine-rich repeats containing G protein-coupled receptor; CTCK, C-terminal cystine knot; WGS, whole-genome sequencing; Burs $\alpha$ , bursicon  $\alpha$  subunit; Burs $\beta$ , bursicon  $\beta$  subunit

**A**

<i>bm</i> Bursa $\alpha$	MSVLNTFLVIVALILCYVNDPVTG-----HEVQLPPGTFKFFC	38
<i>ag</i> Bursa $\alpha$	MKSTFLVVLELAFFLLPGRVLYA-----QKDESDGGSHYSS	36
<i>dm</i> Bursa $\alpha$	MLRHLLRHENNKVFLILLYCVLVSILKLCCTA-----QPDSVAATDNDITHLG	49
<i>am</i> Burs	MKENFSIMFIHSIFLILIIIFIYSNETIA-----	28
<i>dm</i> Bursa $\beta$	MHVQELLFVAAILVPQCLRA-----LRYS	24
<i>ag</i> Bursa $\beta$	MCNSVRTALAASNCCSIVLCCVLLLTTLTVAVTA-----	35
<i>bm</i> Bursa $\beta$	MNIMITKIFFLVQLFYIVVSKSSA-----	24
<i>bm</i> Bursa $\alpha$	-----	38
<i>ag</i> Bursa $\alpha$	-----	36
<i>dm</i> Bursa $\alpha$	-----	49
<i>am</i> Burs	-----Q-VTDDENCETLQSEVHITKDEYDEIGRLKRTCSGDISV	66
<i>dm</i> Bursa $\beta$	-----Q-GTGDENCETLKSEIHLIKEEFDELGRMQRTCNADVIV	62
<i>ag</i> Bursa $\beta$	-----QHNQADETCETLPSEIHLIKEEYDELGRLYRTCNGDVTIV	74
<i>bm</i> Bursa $\beta$	-----EENCETVASEVHVTKEEYDEMGRLLRSCSGEVSIV	58
<i>bm</i> Bursa $\alpha$	-----	38
<i>ag</i> Bursa $\alpha$	-----	36
<i>dm</i> Bursa $\alpha$	-----	49
<i>am</i> Burs	TKCEGFCNSQVQPSVASTTGFSKECYCCRESYLKERHITLHHCYDADGIKLMNEENGVM	126
<i>dm</i> Bursa $\beta$	NKCEGLCNSQVQPSVITPTGFLKECYCCRESFLKEKVITLTHCYDPDGTRLTSPMGSM	122
<i>ag</i> Bursa $\beta$	NKCEGKCNQVQPSVITATGFLKECYCCRESFLRERQLQLTHCYDPDGVVMTDHEATME	134
<i>bm</i> Bursa $\beta$	NKCEGMCNSQVHPSISSPTGFQKECFCCREKFLRERLVTLTHCYDPDGIREFEENALME	118
<i>pa</i> Burs	-----ESFLR-----LTQEGQASME	
<i>bm</i> Bursa $\alpha$	-----QECQMTAVIHVLKHRGCKPKAIPSFACIGKCTSYVQVSGSK	79
<i>ag</i> Bursa $\alpha$	-----DDCQVTPVIHVLQYPGCVPKPIPSFACIGRCASYIQVSGSK	77
<i>dm</i> Bursa $\alpha$	-----DDCQVTPVIHVLQYPGCVPKPIPSFACVGRASYIQVSGSK	90
<i>am</i> Burs	IKIREPVECKCIKCGV--DECQATPVIHFLOYPGCVPKPIPSYACRGRCSYLVQVSGSK	183
<i>dm</i> Bursa $\beta$	IRLREPTECKCFKCGDFTR-----	141
<i>ag</i> Bursa $\beta$	IRLKEPVDCKCFKCGEMVR-----	153
<i>bm</i> Bursa $\beta$	VRLREPDECECYKCGDFSR-----	137
<i>pa</i> Burs	VK-----DGSSYLQVSGSK	
<i>bm</i> Bursa $\alpha$	IWQMERTCNCCQESGEREATVVLFCPDAQNEEKFRKRVSTKAPLQCMCRPCGSIEESSII	139
<i>ag</i> Bursa $\alpha$	IWQMERSCMCCQESGEREASVSLFCPKAKNGEKKFRKRVSTKAPLECMCRPCTGIEDANVI	137
<i>dm</i> Bursa $\alpha$	IWQMERSCMCCQESGEREAAVSLFCPKVKPGERKFKKVLTKAPLECMCRPCTSIEESGII	150
<i>am</i> Burs	IWQMERSCMCCQESGEREASVSLFCPRAKPGKFKFRKVITKAPLECMCRPCTSVEEYAI	243
<i>pa</i> Burs	IWQMER-----EASVNNV	
<i>bm</i> Bursa $\alpha$	PQEVAGYSEEGPLYN--HFRKS--L	160
<i>ag</i> Bursa $\alpha$	PQELTSFADEGTLTGY--FQKSHYKSIE	163
<i>dm</i> Bursa $\alpha$	PQEIAGYSDEGPLNN--HFRRIA-LQ	173
<i>am</i> Burs	PQEIAGFADEGPFPTSAHFRRSSDLQ	269

**B**

<i>am</i> Bursa $\beta$	30	VTDDENCETLQSEVHITKDEYDEIGRLKRTCSGD-LSVTKCEGFCNSQVQPSVAST	84
<i>dm</i> Bursa $\beta$	26	GTGDENCETLKSEIHLIKEEFDELGRMQRTCNADVI-VNKCEGLCNSQVQPSVITP	80
<i>dm</i> Bursa $\alpha$	50	---DDCQVTP-----VIHVLQYPGCVPKPIPSFACVGRASYIQVSGSKI	91
<i>am</i> Bursa	141	--GVDECQATP-----VIHFLOYPGCVPKPIPSYACRGRCSYLVQVSGSKI	184
CTCK	1	KSPVVRQTITYNGCTSVVVKNAFCGKCGSAVMYSIEAQ	38
<i>am</i> Bursa $\beta$	85	TGFSKECYCCRESYLKERHITLHHCYDADGIKLMNEENGVM	140
<i>dm</i> Bursa $\beta$	81	TGFLKECYCCRESFLKEKVITLTHCYDPDGTRLTSPMGSM	136
<i>dm</i> Bursa $\alpha$	92	WQMERSCMCCQESGEREAAVSLF-CPKVKPG-----ERKFKKVLTKAPLECMCRPC	141
<i>am</i> Bursa	185	WQMERSCMCCQESGEREASVSLF-CPRAKPG-----EKKFRKVITKAPLECMCRPC	234
CTCK	39	DVDHS-GSCCQPHKTKTRQVRLR-CPDGST-VTKT-----VMHIECGCEPC	82

Fig. 1. Alignments of insect bursicon proteins. (A) CG13419 and CG15284 orthologs (called Bursa $\alpha$  and Bursa $\beta$ , respectively) from mosquito (*ag*: *Anopheles gambiae*), silkworm (*bm*: *Bombyx mori*) and fruit fly (*dm*: *Drosophila melanogaster*) were aligned along the bee (*am*: *Apis mellifera*) single-chain multidomain bursicon. The tryptic peptides identified in bursicon preparations from the cockroach (*pa*: *Periplaneta americana*), a heterometabolic insect, are also aligned. Conserved Bursa $\alpha$  and Bursa $\beta$  sequences are shaded in grey and yellow, respectively. Residues identical among 3, 4, or 5 species are dark shaded, while residues identical between two species are pale shaded. Silkworm Bursa $\alpha$  (BN000691) and Bursa $\beta$  (BN000690), mosquito Bursa $\beta$  (BN000689) and bee single-chain bursicon (BN000692) were manually predicted on available WGS DNA sequences (see Section 2 for details). *Drosophila* CG13419 and CG15284 cDNAs were cloned, sequenced and annotated in EMBL/GenBank/DBJ databases (Accession Nos. AJ862523 and AJ862524, respectively). The putative signal peptides are shown at the left side on the top of the figure. (B) Alignment of *Drosophila* Bursa $\alpha$  and Bursa $\beta$  proteins, together with the Bursa $\alpha$  and Bursa $\beta$  domains from bee single-chain bursicon, and a consensus cystine knot-like domain (CTCK: C-terminal cystine knot) [30]. Conserved cysteine residues are highlighted. Other conserved residues are shaded in blue.

In the present study we have used a combination of comparative genomic, biological and biochemical approaches to identify bioactive bursicon as a cystine knot heterodimeric protein and the natural agonist of DLGR2.

## 2. Methods

### 2.1. Similarity searches and protein alignments

Similarity searches of insect genomes and proteins were made on the NCBI platform ([www.ncbi.nih.gov](http://www.ncbi.nih.gov)) by TBLASTN or BLASTP, respectively [21]. Multiple protein alignments were performed with CLUSTAL ([www.clustalw.genome.jp](http://www.clustalw.genome.jp)) using default settings.

### 2.2. Protein predictions on WGS (whole-genome sequencing) DNA sequences

The bee, mosquito and silkworm genomes were analysed for the presence of *Drosophila* CG13419 or CG15284 orthologs with TBLASTN (*Anopheles* CG13419-like sequence was already present in the database; Accession No. AY735443). Because the N-termini are not conserved among species, the putative signal peptides could not be identified in the TBLASTN alignments. In consequence, each locus was individually screened for the presence of potential signal peptides [22] located in frame in the same exon with CG13419 or CG15284 homologous sequences (ORF finder tool at the NCBI, [www.ncbi.nih.gov/gorf](http://www.ncbi.nih.gov/gorf); SignalP 3.0 Server; [www.cbs.dtu.dk/services/SignalP/](http://www.cbs.dtu.dk/services/SignalP/)). In the case of the silkworm CG13419 ortholog, the putative signal peptide was identified in a predicted separate exon, located 56 base pairs upstream from the starting point of CG13419-like segment. Exon prediction was performed at [www.fruitfly.org/seq\\_tools/splince.html](http://www.fruitfly.org/seq_tools/splince.html). Bee, silkworm and mosquito predicted sequences were submitted to EMBL/GenBank/DBJ databases. For accession numbers see legend of Fig. 1.

### 2.3. PCR, cloning and sequencing

*Drosophila* cDNAs, normalized to the housekeeping *Rp49* (Ribosomal protein 49) cDNA, derived from different developmental stages and tissues ('Rapid-Scan™ Gene expression panel', Origene Technologies, Rockville, MD) were used as template to amplify bursicon  $\alpha$  subunit (Burs $\alpha$ ) and bursicon $\beta$  subunit (Burs $\beta$ ) cDNAs by means of PCR with gene-specific primers [Burs $\alpha$  (cg13419): fwd gcggatcctcg-cagtcgggcccagcagatg; rev gcgaattcagtcgtaaactgatgggggaatcta]. They contain a 5' *Bam*HI and *Eco*RI restriction site (underlined), respectively. Annealing temperature 55 °C. Burs $\beta$  (cg15284): fwd atcgatgcc-aggaactgctc; rev ttaactgtgaaatgccaca. Annealing temperature 57 °C]. Both reactions were performed for 35 cycles. The resulting products were analysed by agarose gel electrophoresis, purified, cloned into pCR4-TOPO (Invitrogen, Carlsbad, CA) and sequenced (ABI Prism 310, Applied Biosystems, Foster City, CA). ORFs coding for the bursicon subunits were further subcloned into pcDNA3.1 (Invitrogen) for transfection of COS-7 cells.

### 2.4. Preparation of conditioned media

COS-7 cells growing in 35 mm culture dishes were transfected with 1  $\mu$ g of plasmid DNA using the DEAE-dextran method [23]. The day after transfection, the culture medium (DMEM containing 10% Fetal Bovine Serum (FBS)) was replaced by the same medium with 1% FBS. Twenty-four hours later, the medium was recovered, centrifuged at 2000  $\times$  g to eliminate cellular debris, immediately frozen in liquid nitrogen and stored at -20 °C until use. For dose-response curves and gel filtration chromatography, conditioned medium was concentrated using ultrafiltration devices with a 10 kDa MW cut-off (Amicon Ultra Centrifugal Filters, Millipore Corporation).

### 2.5. Stimulation of cAMP accumulation

*Drosophila* DLGR1, DLGR2 [19,28] and human hLGR4 and hLGR5 cloned in pcDNA3 were transfected in COS-7 cells in the same way as described above. Forty-eight hours after transfection, expression of the various receptors was assayed by flow immunocytometry, using a panel of monoclonal antibodies (not shown). Cells were

washed once with Krebs–Ringer–HEPES (KRH) buffer and incubated at 37 °C for 30 min. Buffer was then replaced by 1 ml of fresh KRH supplemented with 10% of conditioned medium and 25  $\mu$ M rolipram (a cAMP phosphodiesterase inhibitor, Laboratoire Logeais, Paris, France). After a 30 min incubation, the level of intracellular cAMP was measured by RIA as described previously [23].

### 2.6. Gel filtration chromatography

One millilitre of 20 $\times$  concentrated culture medium from cells co-expressing recombinant CG13419 and CG15284 proteins was applied onto a gel filtration column (HILoad™ 16/60 Superdex™ 75 prep Grade, Amersham Biosciences, Freiburg, Germany). Calibrations and runs were made in Na<sub>3</sub>(PO<sub>4</sub>) 20 mM/NaCl 200 mM buffer. Eluates corresponding to sizes between 70 and 10 kDa were collected in 2 ml fractions and frozen immediately in liquid nitrogen. One hundred, or 5  $\mu$ l of each fraction were assayed for cAMP stimulating activity on COS-7 cells expressing DLGR2, or in the bursicon bioassay, respectively.

### 2.7. Bursicon bioassay

Bursicon bioactivity was determined by means of the flesh fly cuticle tanning bioassay [2]. Flies (*Neobellieria bullata*) were decapitated (or neck-ligated) after eclosion and kept at room temperature for 1.5 h in a humidified petri dish. This initial incubation period allowed for a pre-selection of flies (the vast majority) that did not display any endogenous bursicon activity after decapitation (or neck-ligation). Flies that remained pale were injected with 5  $\mu$ l of solution (saline, CNS extract, 1 mM 8Br-cAMP, conditioned culture media or chromatographic fractions) and cuticular tanning was evaluated 3 h post-injection. Qualitative evaluation of tanning was as follows: flies do not tan: "Defective tanning"; only the abdomen tans: "Intermediate tanning"; both the abdomen and the thorax tan: "Full Tanning". The CNS extract was prepared from freshly eclosed flesh flies. CNS were dissected and homogenized in saline solution (127.7 mM NaCl, 4.7 mM KCl) and 0.25 CNS-equivalents were injected into each fly.

## 3. Results and discussion

### 3.1. *Drosophila* CG13419 has no Bursicon bioactivity and does not activate DLGR2

With the aim of establishing a direct experimental link between the cystine knot protein described by Dewey et al. [17] (CG13419, hereafter Burs $\alpha$ ) and bursicon, we transfected COS-7 cells with Burs $\alpha$  cDNA inserted in an expression plasmid. The conditioned medium was tested for bursicon bioactivity in freshly eclosed, neck-ligated *N. bullata* (see Fig. 2C and Section 2), and for its ability to stimulate cAMP production in COS-7 cells expressing DLGR2 (Fig. 2A). Contrary to what was suggested on the basis of genetic data by Dewey et al. [17], Burs $\alpha$  was devoid of bursicon bioactivity and incapable of activating DLGR2. Together with the evidence from loss-of-function mutations of Burs $\alpha$  [17], these results indicated that a piece of information was lacking in the bursicon puzzle, most probably a dimerisation partner for Burs $\alpha$ .

### 3.2. Identification of a new Bursicon candidate

Honegger et al. [16] reported a set of peptide sequences from purified bursicon preparations of the cockroach *P. americana*. It is from some of these peptides that CG13419 (Burs $\alpha$ ) was identified in the *Drosophila* genome [17].

Using the sequence of one of these peptides which was initially considered as a contaminant in the bursicon preparation (peptide sequence: ESFLR, see Fig. 1), we found a new ORF in *Drosophila* encoding a 15 kDa cystine knot protein




Fig. 2. Biological activities of *Drosophila bursicon* proteins. (A) Agonist activity on DLGR2. Conditioned media from control COS-7 cells (Basal) or cells expressing Burs $\alpha$ , Burs $\beta$ , or both proteins together (Burs $\alpha\beta$ ), were evaluated for their ability to induce cAMP accumulation in COS-7 cells transfected with *Drosophila* (DLGR1 and DLGR2) and human (hLGR4 and hLGR5) receptors (mock: cells transfected with empty plasmid). (B) Concentration-effect curve. Conditioned media were either diluted or concentrated (from 1/300 dilution to 30 $\times$  concentration) to obtain serial concentrations of DLGR2 agonist activity. Error bars represent the range of duplicate determinations. (C) Bursicon bioassay. Saline, 8Br-cAMP, CNS extract or conditioned media from COS-7 cells transfected with empty plasmid (control medium) or from cells expressing Burs $\alpha$ , Burs $\beta$ , or both (Burs $\alpha\beta$ ) were injected into decapitated or neck-ligated, freshly eclosed flesh flies (*Neobellieria bullata*) to analyse their capacity to stimulate cuticular tanning. The number of flies tested per condition is given in parentheses. White, grey and black bars represent the percentage of flies showing defective, intermediate (abdomen) or full (thorax + abdomen) tanning, respectively. At the right side, photographs of flies representative of each tanning pattern.

(CG15284, hereafter called Burs $\beta$ ). We identified the corresponding orthologous sequences in mosquito (*A. gambiae*), silkworm (*Bombyx mori*) and honey bee (*A. mellifera*) genomes, and manually predicted the corresponding ORFs (Fig. 1A). The encoded proteins are 72%, 63% and 65% identical to *Drosophila* Burs $\beta$ , respectively.

### 3.3. In the honey bee Burs $\alpha$ and Burs $\beta$ are encoded in a single ORF

Whereas the fly, mosquito and silkworm CG15284-like (Burs $\beta$ ) sequences are predicted to be of similar length (141, 153 and 137 residues, respectively), the bee sequence constitutes the aminoterminal portion of a larger conceptual protein (269 aminoacids, Fig. 1A). It harbours a second cystine knot domain at its C-terminus, suggesting that it originated from the intragenic duplication of an ancient motif. In Fig. 1B, we show both cystine knot domains of the bee protein and the two *Drosophila* Burs $\alpha$  and Burs $\beta$  proteins, aligned with the cystine knot consensus sequence (CTCK: C-terminal cystine knot). Unexpectedly, detailed inspection of this second cystine knot domain, revealed that it corresponds to the bee ortholog of Burs $\alpha$  (CG13419) (Fig. 1A). In the other available insect species, Burs $\alpha$  and Burs $\beta$  are clearly encoded in distinct loci. Considering that a multidomain protein, encoded in a single gene in one species, may correspond to a multimeric protein, with each subunit encoded by separate genes, in another species [24–26], we hypothesized that *Drosophila* bioactive bursicon would be a heterodimer made of CG13419 and CG15284 subunits.

### 3.4. Co-expression of recombinant *Drosophila* Burs $\alpha$ and Burs $\beta$ generates bursicon bioactivity and DLGR2 stimulating activity

To test this hypothesis, Burs $\beta$  cDNA was cloned in the expression vector pcDNA3.1 and transfected in COS-7 cells, either alone or together with Burs $\alpha$  plasmid. Conditioned media were prepared from cultures of transfected cells and tested for their ability to stimulate cAMP accumulation in cells expressing DLGR2. Cells transfected with the cDNA of vertebrate or invertebrate orphan receptors belonging to the same subfamily of GPCRs (hLGR4, hLGR5 and DLGR1) [27,28] were used as controls. As can be seen in Fig. 2A, an 8-fold increase in cAMP production was induced in DLGR2-expressing cells, only when stimulated with conditioned medium from cells co-transfected with Burs $\alpha$  and Burs $\beta$  cDNAs. Stimulation with serial dilutions of the medium generated the expected sigmoid concentration–action curve (Fig. 2B). The conditioned media from cells transfected with Burs $\alpha$  or Burs $\beta$  plasmids alone had no effect (Fig. 2A). No stimulation was observed neither with a mixture made of equal amounts of media from cells transfected separately with Burs $\alpha$  or Burs $\beta$  (not shown). This indicates that, in order to achieve DLGR2 agonistic activity, Burs $\alpha$  and Burs $\beta$  need to be co-expressed in the same cell. DLGR1, a *Drosophila* receptor for a still unknown ligand [28], displayed strong constitutive activity [20] but was not stimulated. Neither were the human hLGR4 and hLGR5 receptors (Fig. 2A).

The identity between bursicon and the material present in the medium of cells co-transfected with *Drosophila* Burs $\alpha$




Fig. 3. Molecular weight estimation. Twenty times concentrated conditioned medium, produced in COS-7 cells co-transfected with Burs $\alpha$  and Burs $\beta$  cDNAs, was size fractionated by gel filtration chromatography. Eluted fractions between 70 and 10 kDa were analysed for their ability to stimulate cAMP accumulation in COS-7 cells expressing DLGR2. Grey bars represent the cAMP value after subtraction of the cAMP level of non-stimulated DLGR2 cells. The chromatogram (blue line) corresponds to 280 nm absorbance curve of the proteins used as molecular weight markers (67 kDa: Albumin; 43 kDa: Ovalbumin; 35 kDa:  $\beta$ -Lactoglobulin; 25 kDa: Chymotrypsinogen; 13.7 kDa: Ribonuclease A). Numbers in abscissa indicate the volume eluted from the column (ml). Error bars represent the standard error of four independent cAMP values obtained from each fraction. The lower panel (neck-ligation bioassays) shows the effects of injecting the fractions in neck-ligated flies (–, no tanning response; +, intermediate tanning; ++, full tanning;  $n = 3$  per fraction).

and Burs $\beta$  cDNAs was established by microinjection in freshly eclosed, decapitated or neck-ligated *Neobellieria (Sarcophaga) bullata* flesh flies (Fig. 2C). Strong tanning of the cuticle was observed 3 h post-injection in flies injected with medium from COS-7 cells co-expressing Burs $\alpha$  and Burs $\beta$ . This response was also observed in flies injected with CNS extract derived from freshly eclosed flesh flies or with 8Br-cAMP, a membrane permeable cyclic AMP analog previously shown to mimic bursicon activity [13,14]. However, no bursicon activity was observed in flies injected with saline, control medium or media derived from cells transfected with Burs $\alpha$  or Burs $\beta$  cDNA alone.

### 3.5. *Drosophila Bursicon* is a heterodimer made of Burs $\alpha$ and Burs $\beta$ cystine knot subunits

The molecular weight of the bioactive molecule present in the conditioned medium was estimated by size-exclusion chromatography (Fig. 3). Eluted fractions were evaluated for bursicon bioactivity and their capacity to stimulate cAMP accumulation in DLGR2 expressing cells. Both activities co-eluted in the 30–35 kDa region of the chromatogram (Fig. 3). These results are in agreement with the expected molecular weights of Burs $\alpha$  and Burs $\beta$  protomers (15.5 and 13.8 kDa, after signal peptide removal, respectively), with the size reported for bioactive hormone [15], and demonstrate the heterodimeric nature of bursicon.

### 3.6. Burs $\alpha$ and Burs $\beta$ mRNAs are expressed at the same developmental stages

The expression of Burs $\alpha$  and Burs $\beta$  transcripts in *Drosophila melanogaster* was explored by RT-PCR at various stages of development. Both transcripts were found in embryonic, larval and pupal stages, but not in the adult of the fruit fly (Fig. 4). This pattern of expression of Burs $\alpha$  and Burs $\beta$  transcripts during development correlates with the one previously reported for DLGR2 [19].




Fig. 4. Expression of Burs $\alpha$  and Burs $\beta$  transcripts in different developmental stages of the fruit fly, *Drosophila melanogaster*, as shown by PCR on normalized cDNA samples (see Section 2). Lanes 1–4: embryonic stages (0–4/4–8/8–12/12–24 h), lanes 5–7: larval stages 1–3, lane 8: pupal stage, lanes 9–10: adult head (males and females), lanes 11–12: adult body (males and females), M: marker (100 bp ladder).

## 4. Conclusion

We have demonstrated that the molting neurohormone bursicon is a heterodimeric protein made of two cystine knot subunits, and the natural agonist of the previous orphan G protein-coupled receptor DLGR2. A similar situation exists in vertebrates, where the pituitary hormones FSH, LH and TSH are also heterodimeric cystine knot proteins acting on their respective leucine-rich repeats containing receptors. The orphan GPCRs LGR4, LGR5 and LGR6 are putative orthologs of DLGR2 in vertebrates [29]. The knowledge of bursicon structure might help in the identification of candidate agonists for these orphan receptors, amongst the large family of cystine knot proteins. The two bursicon protomers, Burs $\alpha$  and Burs $\beta$ , are encoded by separate genes in *Drosophila*, *Anopheles* and *Bombyx*, but they are fused in a single gene in *Apis*. The sequence similarity between Burs $\alpha$  and Burs $\beta$  and their presence as a predicted single-chain protein in *A. mellifera* is a strong indication that they originated by duplication of (or in) an ancestral gene encoding a cystine knot protein.

From a practical point of view, our study makes it possible to develop screening assays for low molecular weight leucine-rich repeats containing G protein-coupled receptor 2 (LGR2) antagonists. These might qualify as effective novel agents for insect pest management.

**Acknowledgements:** We thank Katrien Backers and David Comuni for help with gel filtration chromatography, Sofie Van Soest for DNA sequencing, Roger Jonckers for rearing of flies and Julie Puttemans for photography. Supported by the 'Belgian Programme on Interuniversity Poles of Attraction' (IUAP/PAI P5/30) initiated by the Belgian State, Prime Minister's Office Science Policy Programming. Also supported by grants from the 'Fonds de la Recherche Scientifique Médicale', 'Fonds National de la Recherche Scientifique', Francqui Foundation, 'Fonds voor Wetenschappelijk Onderzoek' (FWO), Lundbeck Foundation, Novo Nordisk Foundation, and Carlsberg Foundation. Tom Van Loy obtained a PhD fellowship from the 'Instituut voor de aanmoediging van Innovatie door Wetenschap en Technologie in Vlaanderen' (IWT). Jeroen Poels obtained a post-doctoral fellowship from the 'Onderzoeksfonds KULeuven'.

## References

- [1] Fraenkel, G. and Hsiao, C. (1962) Hormonal and nervous control of tanning in the fly. *Science* 138, 27–29.
- [2] Fraenkel, G. and Hsiao, C. (1963) Tanning in the adult fly: A new function of neurosecretion in the brain. *Science* 141, 1057–1058.
- [3] Cottrell, C.B. (1962) The imaginal ecdysis of blowflies. *J. Exp. Biol.* 39, 413–431.
- [4] Fraenkel, G., Hsiao, C. and Seligman, M. (1966) Properties of bursicon: An insect protein hormone that controls cuticular tanning. *Science* 151, 91–93.
- [5] Horodyski, F.M., Riddiford, L.M. and Truman, J.W. (1989) Isolation and expression of the eclosion hormone gene from the tobacco hornworm, *Manduca sexta*. *Proc. Natl. Acad. Sci. USA* 86, 8123–8127.
- [6] Zitnan, D., Kingan, T.G., Hermesman, J.L. and Adams, M.E. (1996) Identification of ecdysis-triggering hormone from an epitracheal endocrine system. *Science* 271, 88–91.
- [7] Baker, J.D., McNabb, S.L. and Truman, J.W. (1999) The hormonal coordination of behavior and physiology at adult ecdysis in *Drosophila melanogaster*. *J. Exp. Biol.* 202 (Pt 21), 3037–3048.
- [8] Fuse, M. and Truman, J.W. (2002) Modulation of ecdysis in the moth *Manduca sexta*: The roles of the suboesophageal and thoracic ganglia. *J. Exp. Biol.* 205, 1047–1058.
- [9] Park, J.H., Schroeder, A.J., Helfrich-Forster, C., Jackson, F.R. and Ewer, J. (2003) Targeted ablation of CCAP neuropeptide-containing neurons of *Drosophila* causes specific defects in execution and circadian timing of ecdysis behavior. *Development* 130, 2645–2656.
- [10] Park, Y., Filippov, V., Gill, S.S. and Adams, M.E. (2002) Deletion of the ecdysis-triggering hormone gene leads to lethal ecdysis deficiency. *Development* 129, 493–503.
- [11] Clark, A.C., del Campo, M.L. and Ewer, J. (2004) Neuroendocrine control of larval ecdysis behavior in *Drosophila*: Complex regulation by partially redundant neuropeptides. *J. Neurosci.* 24, 4283–4292.
- [12] Fraenkel, G., Blechl, A., Blechl, J., Herman, P. and Seligman, M.I. (1977) 3':5'-cyclic AMP and hormonal control of puparium formation in the fleshfly *Sarcophaga bullata*. *Proc. Natl. Acad. Sci. USA* 74, 2182–2186.
- [13] Baker, J.D. and Truman, J.W. (2002) Mutations in the *Drosophila* glycoprotein hormone receptor, rickets, eliminate neuropeptide-induced tanning and selectively block a stereotyped behavioral program. *J. Exp. Biol.* 205, 2555–2565.
- [14] Kimura, K., Kodama, A., Hayasaka, Y. and Ohta, T. (2004) Activation of the cAMP/PKA signaling pathway is required for post-ecdysial cell death in wing epidermal cells of *Drosophila melanogaster*. *Development* 131, 1597–1606.
- [15] Kostron, B., Market, D., Kellermann, J., Carter, C.E. and Honegger, H.W. (1999) Antisera against *Periplaneta americana* Cu,Zn-superoxide dismutase (SOD): Separation of the neurohormone bursicon from SOD, and immunodetection of SOD in the central nervous system. *Insect Biochem. Mol. Biol.* 29, 861–871.
- [16] Honegger, H.W., Market, D., Pierce, L.A., Dewey, E.M., Kostron, B., Wilson, M., Choi, D., Klukas, K.A. and Mesce, K.A. (2002) Cellular localization of bursicon using antisera against partial peptide sequences of this insect cuticle-sclerotizing neurohormone. *J. Comp. Neurol.* 452, 163–177.
- [17] Dewey, E.M., McNabb, S.L., Ewer, J., Kuo, G.R., Takanishi, C.L., Truman, J.W. and Honegger, H.W. (2004) Identification of the gene encoding bursicon, an insect neuropeptide responsible for cuticle sclerotization and wing spreading. *Curr. Biol.* 14, 1208–1213.
- [18] Riehle, M.A., Garczynski, S.F., Crim, J.W., Hill, C.A. and Brown, M.R. (2002) Neuropeptides and peptide hormones in *Anopheles gambiae*. *Science* 298, 172–175.
- [19] Eriksen, K.K., Hauser, F., Schiott, M., Pedersen, K.M., Sondergaard, L. and Grimmelikhuijzen, C.J. (2000) Molecular cloning, genomic organization, developmental regulation, and a knock-out mutant of a novel leu-rich repeats-containing G protein-coupled receptor (DLGR-2) from *Drosophila melanogaster*. *Genome Res.* 10, 924–938.
- [20] Nishi, S., Hsu, S.Y., Zell, K. and Hsueh, A.J. (2000) Characterization of two fly LGR (leucine-rich repeat-containing, G protein-coupled receptor) proteins homologous to vertebrate glycoprotein hormone receptors: Constitutive activation of wild-type fly LGR1 but not LGR2 in transfected mammalian cells. *Endocrinology* 141, 4081–4090.
- [21] Altschul, S.F., Madden, T.L., Schaffer, A.A., Zhang, J., Zhang, Z., Miller, W. and Lipman, D.J. (1997) Gapped BLAST and PSI-BLAST: A new generation of protein database search programs. *Nucleic Acids Res.* 25, 3389–3402.
- [22] Bendtsen, J.D., Nielsen, H., von Heijne, G. and Brunak, S. (2004) Improved prediction of signal peptides: SignalP 3.0. *J. Mol. Biol.* 340, 783–795.
- [23] Vlaeminck, V., Ho, S.C., Rodien, P., Vassart, G. and Costagliola, S. (2002) Activation of the cAMP pathway by the TSH receptor involves switching of the ectodomain from a tethered inverse agonist to an agonist. *Mol. Endocrinol.* 16, 736–746.
- [24] Enright, A.J., Iliopoulos, I., Kyrpides, N.C. and Ouzounis, C.A. (1999) Protein interaction maps for complete genomes based on gene fusion events. *Nature* 402, 86–90.
- [25] Marcotte, E.M., Pellegrini, M., Ng, H.L., Rice, D.W., Yeates, T.O. and Eisenberg, D. (1999) Detecting protein function and protein–protein interactions from genome sequences. *Science* 285, 751–753.
- [26] Sali, A. (1999) Functional links between proteins. *Nature* 402, 25–26.
- [27] Hsu, S.Y., Liang, S.G. and Hsueh, A.J. (1998) Characterization of two LGR genes homologous to gonadotropin and thyrotropin receptors with extracellular leucine-rich repeats and a G protein-coupled, seven-transmembrane region. *Mol. Endocrinol.* 12, 1830–1845.
- [28] Hauser, F., Nothacker, H.P. and Grimmelikhuijzen, C.J. (1997) Molecular cloning, genomic organization, and developmental regulation of a novel receptor from *Drosophila melanogaster* structurally related to members of the thyroid-stimulating hormone, follicle-stimulating hormone, luteinizing hormone/choriogonadotropin receptor family from mammals. *J. Biol. Chem.* 272, 1002–1010.
- [29] Hsu, S.Y., Kudo, M., Chen, T., Nakabayashi, K., Bhalla, A., van der Spek, P.J., van Duin, M. and Hsueh, A.J. (2000) The three subfamilies of leucine-rich repeat-containing G protein-coupled receptors (LGR): Identification of LGR6 and LGR7 and the signaling mechanism for LGR7. *Mol. Endocrinol.* 14, 1257–1271.
- [30] Marchler-Bauer, A., Anderson, J.B., DeWeese-Scott, C., Fedorova, N.D., Geer, L.Y., He, S., Hurwitz, D.I., Jackson, J.D., Jacobs, A.R., Lanczycki, C.J., Liebert, C.A., Liu, C., Madej, T., Marchler, G.H., Mazumder, R., Nikolskaya, A.N., Panchenko, A.R., Rao, B.S., Shoemaker, B.A., Simonyan, V., Song, J.S., Thiessen, P.A., Vasudevan, S., Wang, Y., Yamashita, R.A., Yin, J.J. and Bryant, S.H. (2003) CDD: A curated Entrez database of conserved domain alignment. *Nucleic Acids Res.* 31, 383–387.
- [31] Luo, C.W., Dewey, E.M., Sudo, S., Ewer, J., Hsu, S.Y., Honegger, H.W. and Hsueh, A.J. (2005) Bursicon, the insect cuticle-hardening hormone, is a heterodimeric cystine knot protein that activates G protein-coupled receptor LGR2. *Proc. Natl. Acad. Sci. USA* 102, 2820–2825.