



UNIVERSITY OF CALIFORNIA PRESS
JOURNALS + DIGITAL PUBLISHING



Global Decline of and Threats to *Aegagropila linnaei*, with Special Reference to the Lake Ball Habit

Author(s): Christian Boedeker, Anja Eggert, Anne Immers, Erik Smets

Reviewed work(s):

Source: *BioScience*, Vol. 60, No. 3 (March 2010), pp. 187-198

Published by: [University of California Press](#) on behalf of the [American Institute of Biological Sciences](#)

Stable URL: <http://www.jstor.org/stable/10.1525/bio.2010.60.3.5>

Accessed: 11/01/2012 09:04

Your use of the JSTOR archive indicates your acceptance of the Terms & Conditions of Use, available at <http://www.jstor.org/page/info/about/policies/terms.jsp>

JSTOR is a not-for-profit service that helps scholars, researchers, and students discover, use, and build upon a wide range of content in a trusted digital archive. We use information technology and tools to increase productivity and facilitate new forms of scholarship. For more information about JSTOR, please contact support@jstor.org.



University of California Press and American Institute of Biological Sciences are collaborating with JSTOR to digitize, preserve and extend access to BioScience.

<http://www.jstor.org>

Global Decline of and Threats to *Aegagropila linnaei*, with Special Reference to the Lake Ball Habit

CHRISTIAN BOEDEKER, ANJA EGGERT, ANNE IMMERS, AND ERIK SMETS

The freshwater macroalga Aegagropila linnaei can occur attached or unattached, but is best known for its free-floating spherical growth form (“lake balls,” “Cladophora balls,” or “marimo”). The worldwide distribution of this species was reconstructed using more than 1200 specimens from 28 herbaria, an extensive literature survey, and field observations. Aegagropila linnaei was found in 283 locations, with the majority located in central and northern Europe. Few records of A. linnaei exist from North America, but it is relatively widespread in Japan. The preferred habitat is inferred to be oligomesotrophic lakes with moderate to high calcium levels. In more than 50% of the known locations, A. linnaei has not been found in the last 30 years, and this decline is assumed to be correlated with changes in trophic conditions in those lakes. The unattached growth forms show a stronger decline than attached populations as a result of the habitat types in which they can occur.

Keywords: aquarium trade, Cladophora balls, eutrophication, freshwater, unattached growth form

The freshwater green alga *Aegagropila linnaei* (Cladophorophyceae, Chlorophyta) is best known for its unattached spherical growth form, also referred to as “marimo,” “lake balls,” or “*Cladophora* balls” (figure 1a). These lake balls can develop under certain hydrographic and topographic conditions either directly from the attached growth form (figure 1b), from floating unattached mats, or from fragmented, older balls (Brand 1902, Kurogi 1980). Individual filaments of *A. linnaei* have a simple morphology of uniseriate branched filaments similar to *Cladophora* species (figure 1c). The filaments can be attached by a holdfast or secondary rhizoids to the substratum, forming low turfs on submerged rocks or bivalves, or can be intermingled and free floating as mats or balls (van den Hoek 1963, Leliaert and Boedeker 2007). The lake balls are radially organized and increase in size by the vegetative growth of the interwoven filaments. The stiff texture of the filaments and the production of secondary rhizoids are features that facilitate entanglement and subsequent ball formation. Water motion is the main shape-giving factor, and balls occur typically in shallow lakes with sandy bottoms. Rolling movements on the sediment and abrasion of apical cells induce radial growth and profuse lateral branching (Acton 1916). Different growth forms can co-occur under different conditions in the same lake (Luther 1951, Kurogi 1980).

The spherical growth form has attained considerable popularity. *Aegagropila linnaei* has been given the status of

a “special natural monument” in Japan (Kurogi 1980), and the lake balls are shown on postage stamps from Japan and Iceland. In Japan, a whole range of marimo merchandise exists, ranging from souvenirs and lucky charms with live specimens to toys and cartoons. *Aegagropila linnaei* balls have also become extremely popular in the aquarium trade in recent years.

In contrast to most other freshwater algae, *A. linnaei* has a restricted distribution, and is found mainly in previously glaciated areas of Europe. In addition, *A. linnaei* has been identified in a range of locations in Japan, whereas only few records are known from North America. In most countries it is regarded as a rare species. Freshwater algae that are not widespread are often restricted in their distribution by specific ecological demand, such as desmids (Coesel et al. 1978) or macroscopic *Nostoc* species (Mollenhauer et al. 1999). On the other hand, a restricted distribution can also be the result of a limited dispersal capacity. Migrating birds are one of the most important dispersal vectors for freshwater algae (Schlichting 1960, Green et al. 2002), transporting desiccation-resistant stages such as akinetes. However, akinetes are basically unknown in *A. linnaei* (Brand 1902, Acton 1916, van den Hoek 1963). Limited dispersal abilities and competition effects have most likely shaped the current distribution of *A. linnaei* more than has the availability of habitats.

Even though the species was initially assumed to occur mainly in eutrophic waters (van den Hoek 1963), eutrophication is



Figure 1. (a) A lake ball, ordered from an online aquarium shop. (b) Attached growth form of *Aegagropila linnaei* from Lake Erken, Sweden. Photograph: Roland Bengtsson. (c) Microscopical characters of *A. linnaei* filaments with typical subterminal and serial insertion of branches. (d) Herbarium box (in botanical Garden and Museum Berlin-Dahlem, Germany) with old specimens (such as *Aegagropila sauteri*) from a lake in Sweden.

mentioned as a threat in several more recent studies reporting declining populations of *A. linnaei*. Partly because of the popularity of the intriguing ball form, this is one of the few freshwater algal species that has been listed as threatened on several red lists of endangered species. *Aegagropila linnaei* is

included in the national red lists of Belarus (MINPRIRODA 2006), Estonia (Lilleleht 1998), Germany (Ludwig and Schnitler 1996), Japan (EAJ 2000), Russia (Noskov 2000), and Sweden (Gärdenfors 2005). In addition, *A. linnaei* is explicitly mentioned in the Ramsar specifications of Lake Akan, Japan

(Moriwake 2005) and Lake Mývatn, Iceland (Gíslason 1992). In the United Kingdom, *A. linnaei* is included the list of rare algae and in the assessments of Important Plant Areas by Plantlife (Brodie et al. 2008).

For several locations, this species is already assumed or known to be extinct. In four out of the five locations in the Netherlands where *A. linnaei* was found 60 years ago, it is now absent, and the unattached forms seem to have disappeared altogether as a result of the effects of eutrophication (Boedeker and Immers 2009). Balls of *A. linnaei* have been extinct in Lake Zeller in Austria since around 1910, most likely due to effects of human activity (Nakazawa 1974), and only the attached filamentous form has been reported since then (Kann and Sauer 1982). The attached growth form is still present in several larger rivers of northern England and Scotland, but the ball-shaped growth form is found only in a few unchanged locations (John 2002). There is only one location in Denmark (Sorø Sø, Sjøland) where *A. linnaei* occurs, although the species had been found previously in more locations there (van den Hoek 1963). In Lake Galenbecker, in northeastern Germany, carp cultivation and intensified agriculture in the 1960s led to eutrophication and extinction of the species (Pankow 1985). In Japan, where the species is still relatively widespread, human activities have led to the destruction of many of *A. linnaei*'s natural habitats. Populations of *A. linnaei* have disappeared from two bays in Lake Akan, most likely as a result of increased mud deposition (Wakana et al. 2006). Population declines (both in population density and distribution area) have been monitored in the Takkobu Marsh, Japan, between 1996 and 2004, and are probably a result of the effects of eutrophication, such as higher nutrient load, accumulating layers of mud and silt on the marsh bottom, shading by phytoplankton blooms, and an increase in the depth limit of freshwater bivalves (Wakana et al. 2005). Declining populations have also been observed in several other small swamp lakes for the same reasons (Wakana et al. 2001). The *Red Data Book of Estonia* lists eutrophication, changes in water hydrology, and dredging as factors that threaten the survival of *A. linnaei* (Lilleleht 1998).

In this study, we used detailed information on the distribution of *A. linnaei* compiled from herbarium specimens (figure 1d), literature data, and field observations to assess whether there is a global trend of declining populations. Habitat preferences with regard to trophic level are inferred, and we discuss possible deleterious effects of eutrophication. Furthermore, we evaluated whether unattached populations, including the ball form, are more threatened than the attached growth form.

Location survey

We surveyed herbarium collections to produce detailed distribution maps of *A. linnaei*. Fifty-nine herbaria responded to our inquiries, of which 28 had collections of *A. linnaei* (for a list of *A. linnaei* synonyms, see appendix S1 in the supplementary material at www.nationaalherbarium.nl).

[nl/supplements/boedeker/BioScience_2010](http://www.nationaalherbarium.nl/supplements/boedeker/BioScience_2010); see appendix S2 for a complete list of herbaria). We borrowed from or visited collections, and identified the samples using a light microscope (Olympus BH2) after resoaking fragments in water. In total, we checked approximately 1200 herbarium specimens of *A. linnaei*, about half of which had location details, and a third of which (about 400 specimens) had a collection date (table 1). Almost all specimens with missing information seem to be about a century old, judging from the general appearance of the vouchers and the handwriting. In addition to the location and collection date, we recorded the growth form wherever possible. The resulting data set was amended by an extensive literature survey and direct inquiries with local and national water-monitoring organizations, or with limnological departments in countries where *A. linnaei* was known or suspected to occur. These inquiries were particularly useful in obtaining information on the existence of extant populations and on past and present trophic levels of the habitats. We reduced this data set to a digest listing all locations (see appendix S3 in the supplementary material at www.nationaalherbarium.nl/supplements/boedeker/BioScience_2010), and georeferenced the locations. Excluded specimens, locations, and synonyms are listed in appendices S1 and S3 online. We created maps using Manifold GIS (geographic information system) software, and displayed data using the World Geodetic 1984 projection. We treated multiple records from the same lake—that is, in different years or in different areas of the same lake—as one location. Almost one-fifth of the locations are from the Baltic Sea, so these locations were grouped together as Baltic proper, western Gulf of Bothnia, eastern Gulf of Bothnia, Gulf of Finland, and Bay of Bothnia, each counting as one location to deflate total location numbers, since individual lakes represent locations in freshwater environments. Records younger than 30 years have been classified as “extant populations” and are listed separately in appendix S4 in the supplementary material (www.nationaalherbarium.nl/supplements/boedeker/BioScience_2010).

Origin of *A. linnaei* balls in the aquarium trade

We contacted more than 20 aquatic plant traders in Belgium, Denmark, Germany, Japan, the Netherlands, Singapore, Ukraine, and the United States to find out about the origin of *A. linnaei* balls for sale in commercial aquarium shops, and to learn about the possible supply from wild stocks.

Results of the location survey

Table 1 shows an overview of the herbarium and literature survey. In total, *A. linnaei* had been recorded from 283 locations (see S3 at www.nationaalherbarium.nl/supplements/boedeker/BioScience_2010), or 233 locations when the list is corrected for the continuous coastline of the Baltic Sea. We sorted the dated herbarium specimens of *A. linnaei* in intervals of 25 years to illustrate the distribution of collections over time (figure 2a, 2b, 2c). The first two time intervals

Table 1. Number of herbarium specimens, locations, and extant populations of *Aegagropila linnaei*.

Type	Number
Herbarium specimens	1200
Herbarium specimens with recorded location	661
Herbarium specimens with recorded date	404
“Ball-shaped” herbarium specimens	310
Locations	
Locations (including literature and field observations)	283
Locations corrected for Baltic Sea ^a	233
Locations with known growth forms ^a	185
Locations with attached growth form ^a	61
Locations with unattached growth form (ball-shaped) ^a	124 (90)
Extant locations (< 30 years)	
Extant locations ^a (percent decline)	101 (57)
Extant locations with attached growth form ^a (percent decline)	33 (46)
Extant locations ^a with unattached/ball-shaped form (percent decline)	51/36 (60)
Top six countries (extant locations)	
Sweden ^a	51 (23)
Germany	33 (8)
Great Britain and Ireland	32 (16)
Japan	21 (20)
Russia ^a	19 (2)
Iceland	10 (9)

a. The Baltic Sea is divided into five areas: Baltic proper, Gulf of Bothnia West, Gulf of Bothnia East, Gulf of Finland, and Bay of Bothnia, each counting as one location.

in each diagram are underrepresented, since about 600 herbarium specimens did not have a date but were clearly very old, as indicated by the dashed lines. Both the number of specimens (figure 2a) and their locations (figure 2b, 2c) show a decrease since the beginning of the last century. This decrease is most pronounced from 1900 onward, and again from 1975 onward. The number of locations represented by herbarium specimens dropped from 32 (1875–1899) to 7 (1975–2007), representing a decline in locations of 78%. The number of locations for ball-shaped specimens dropped from 27 to 5 in the same time interval (figure 2b, 2c). Since herbarium collections reflect not only the occurrence of species at a particular time and place but also the degree of collecting activity, we complemented the herbarium data set with a comprehensive literature survey and field observations, which yielded an additional 152 locations (see S3 in the supplementary material at www.nationaalherbarium.nl/supplements/boedeker/BioScience_2010; combined data shown in figure 2d).

We classified records from less than 30 years ago as “extant populations,” and these are listed by the most recent collection date in the supplementary material S4 (www.nationaalherbarium.nl/supplements/boedeker/

BioScience_2010). *Aegagropila linnaei* has been found in 107 locations since 1976; 101 locations when correcting this number for multiple locations within the same area of the Baltic Sea (table 1; S4 online). Of the 235 total locations where *A. linnaei* had been found originally, it has not been recorded in the last 30 years in about 57% of those locations (figure 2d). All extant populations are indicated in figure 3. Outside Europe, extant populations of *A. linnaei* can be found in a considerable number of locations in both Iceland and Japan (figure 3a). Within Europe, the majority of extant populations exist in Sweden, Great Britain, Ireland, and northeastern Germany (figure 3b)—thus in areas that also had the highest density of original habitats. In the Alps, one of the clusters of original habitats, *A. linnaei* has been reported from only a single location in the last 30 years (Lake Zeller). For the majority of European locations, no recent findings of *A. linnaei* have been made.

Differences in *A. linnaei* population declines have been found for the different growth forms (figure 2d). The presumed loss of locations is less pronounced for the attached growth form (48% decline) than for the unattached forms (59% decline; ball form only, 60% decline).




Figure 2. (a) Number of herbarium specimens of *Aegagropila linnaei* over time. Black bars indicate the total number of herbarium specimens, gray bars indicate the number of herbarium specimens corrected for doublettes. (b) Number of locations of *A. linnaei* over time based on herbarium specimens. (c) Number of herbarium specimens (black bars) and locations based on herbarium specimens (gray bars) of the ball-shaped growth form of *A. linnaei*. (d) Total number of known locations of *A. linnaei* (black bars) and known extant populations (< 30 years; gray bars), based on herbarium specimens, literature, and field data. Dashed lines indicate the large number of very old herbarium specimens that had no date.

A direct comparison of original and extant European locations in relation to the growth forms of *A. linnaei* is given in figure 4. Figure 4a and 4b show all known attached populations (historical and current) and the extant attached populations, respectively. For about 48% of the locations (corrected) with the attached growth form, *A. linnaei* has not been reported in the last 30 years. In figure 4c and 4d, both the original and the extant unattached populations are compared, further broken down into floating and ball-forming populations. Unattached growth forms have been found more commonly in Europe than the attached growth form (figure 4a, 4c). However, when comparing the number of extant ball-forming populations with the number of

historical habitats for this growth form, *A. linnaei* has not been reported in about 60% of those locations in the last 30 years (59% for all unattached growth forms combined).

Origin of *A. linnaei* balls in the aquarium trade: Results

Of more than 20 contacted aquarium shops and aquatic plant trading companies, 13 were willing to share at least some information concerning the origin of the “lake balls” in their product line. Generally, companies were very reluctant to reveal their suppliers, growers, or wholesalers. However, in all instances where information was obtained, Ukraine was given as the country of origin, sometimes specified as Lake Svityaz, or the Shatsk lakes, in northwestern Ukraine.




Figure 3. (a) All known locations of *Aegagropila linnaei* (black dots) and extant populations (< 30 years; white dots). (b) A more detailed map of Europe.

Local Ukrainian sellers of *A. linnaei* balls in Odessa and Kyiv said they receive their material periodically from Lake Svityaz, but claimed that they grow the majority of balls themselves. Most companies grow balls vegetatively from fragments to market size, but receive stocks from Ukraine.

Balls from Lake Svityaz are shipped even to aquatic plant-breeding facilities in Southeast Asia before they are returned to the European market. Japanese aquarium shops sell balls only of European origin. No other natural source of *A. linnaei* balls other than the Shatsk lakes (including Lake




Figure 4. Direct comparisons of all known and extant locations of *Aegagropila linnaei* in Europe, shown separately for attached and unattached growth forms. (a) All known locations of attached populations. (b) Extant locations of attached populations. (c) All known locations of unattached growth forms (black dots: unattached filaments or mats; white dots: balls). (d) Extant locations of unattached growth forms (black dots: unattached filaments or mats; white dots: balls).

Svityaz) was ever mentioned by people in the aquarium plant-trade business.

Evidence for decline of *A. linnaei* populations

Detailed knowledge of the distribution of a species is a prerequisite for (a) assessing the status of that species, (b) monitoring (a decline of) its populations, (c) identifying general threats, and (d) employing most conservation efforts. Typical difficulties in reconstructing the area of distribution are related to incomplete or inaccessible records, patchy local knowledge, and to verifying observations made by third parties. Many organisms, especially among the algae, must be regarded as underrecorded, and are frequently misidentified. This is most likely also the case with *A. linnaei*, despite all efforts to assemble as complete a data set as possible. Even though this species has for a long time intrigued scientists (e.g., Brand 1902, Wesenberg-Lund 1903, Acton 1916, Kindle 1934, Waern 1952), very few reports have provided anything more than scanty lists of findings (Palik 1963, van den Hoek 1963, Pankow 1965, Getzen 1967). All biogeographic reconstructions are hampered by incomplete distribution records, and for most algal species, the knowledge of the global distribution is poor and based on many assumptions. Although some regions of the world are well sampled, such as (western) Europe, the largest parts of the globe are not. The absence of *A. linnaei* in northern North

America and Siberia (figure 3a) could be an artifact of undersampling, despite our efforts to locate records from those areas. Even though it seems unlikely that *A. linnaei* would occur in South America, suitable habitats are potentially abundant in the Andes region, but nobody has looked.

The studies on the extinction or decline of *A. linnaei* cited above are all restricted to a confined area, mostly one individual location. We analyzed as many findings as possible to investigate whether there is a general trend of declining populations. Nevertheless, our record of *A. linnaei* is most likely incomplete, despite the fact that the occurrence of the enigmatic balls in a lake is frequently recorded or commented upon, as illustrated by a wealth of historical scientific and amateur literature. Thus, we assume underrecording is less of a problem for *A. linnaei* than for most other algal species, at least for its unattached growth forms. Even though the observed decline in herbarium specimens does probably also reflect a decline in the submission of voucher specimens, the herbarium collections show the same declining trend as the more comprehensive survey, which also includes data from the literature and communicated field observations (figures 2d, 3). Extant populations are known for less than half of the original locations (see figure 3a). The inferred decline might be affected by another methodological problem; that is, for some regions (e.g., Russia), old literature is available but no or few personal contacts exist, and recent literature is difficult to obtain (see table 1).

We found differences in the degree of decline between attached and unattached growth forms, with the loss of unattached forms being more pronounced (figure 2d). Unattached forms usually float just above the sediment, and therefore occur only in shallow lakes or shallow parts of a deeper lake, where the euphotic zone reaches the bottom. It seems that the stronger decline of unattached growth forms is not caused by a higher sensitivity to environmental changes compared with the attached forms, but rather by a higher threat to their shallow habitats. The attached form most commonly occurs on submerged rocks or freshwater bivalves (Luther 1951, Waern 1952, Pankow and Bolbrinker 1984, Wakana et al. 2001), and has been found in depths of 20 meters (maximum depth 50 meters) in deep lakes (Brand 1902, Kindle 1915), as well as in a number of rivers (Holmes and Whitton 1975, 1977, 1981, Gardavský 1986, Whitton et al. 1998). The attached form might be able to escape greater sedimentation and better cope with decreasing underwater light availability by colonizing substrates closer to the surface, and therefore increasing chances of survival in disturbed habitats. In addition, the ability to survive in deep and steep lakes increases the absolute number of potential habitats for the attached form, and those lakes are generally less sensitive to anthropogenic changes than shallow, lowland lakes are (e.g., Taranu and Gregory-Eaves 2008). The attached form is distributed continuously only in the low-salinity areas of the northern Baltic Sea (figure 2b), where species numbers and competition are low. *Aegagropila linnaei* forms dense carpets on rocks in the Bay and Gulf of Bothnia, and can be the dominant macroalga in areas with a salinity below 4 practical salinity units (Bergström and Bergström 1999). However, the number of recent records from that area indicates a declining population as well (figures 3, 4). The decline of unattached growth forms is most obvious in central Sweden, northeastern Germany, and the subalpine region of the European Alps (figure 4c, 4d).

Eutrophication as the main cause for the decline of *A. linnaei*

Human activities increase and accelerate the external supply of nutrients to aquatic ecosystems worldwide (e.g., Smith et al. 1999, MEA 2005). Freshwater habitats especially face increasing threats from physical alteration, changes in water level and salinity, overexploitation, introduction of non-native species, herbicide and other biocide runoff, airborne pollution, and nutrient loading (MEA 2005, Revenga et al. 2005). Eutrophication of aquatic ecosystems is a common process worldwide and leads to the loss of unique habitats and a reduction in biodiversity (e.g., Bayly and Williams 1973, Smith et al. 1999).

Enrichment of nitrogen and phosphorus loadings selects for fast-growing algae (phytoplankton, macroalgae such as *Ulva* or *Cladophora* species) at the expense of slower-growing species (Duarte 1995), such as some charophytes, for example, or *A. linnaei*. Shallow lakes are especially prone to the regime shift from aquatic macrophytes to phytoplankton dominance.

The decline of several freshwater algal groups has been attributed to eutrophication effects; such groups include macroscopic *Nostoc* species (Mollenhauer 1998, Mollenhauer et al. 1999), desmids (Coesel et al. 1978, Geissler 1988), and charophytes (Geissler 1988, Blindow 1992, Nagasaka et al. 2002). A number of reports on the local decline or extinction of populations of *A. linnaei* mention eutrophication as a responsible factor (Pankow 1985, Wakana et al. 2001, 2005, 2006, Boedeker and Immers 2009).

Our literature study showed that *A. linnaei* occurs in several different lake types, but most typically in shallow, oligomesotrophic, glaciofluvial lakes with reed stands, dense charophyte vegetation, a pH greater than 7, and moderate to high calcium levels. However, the ecological preferences or requirements of *A. linnaei* have never been fully characterized, and conflicting views can be found in the literature with regard to nutrient levels. In a classic volume on European *Cladophora* species, van den Hoek stated that *A. linnaei* (as *Cladophora aegagropila*) “seems to occur only in more or less eutrophic water” (1963). In this study, we collected information on trophic-level changes for a considerable number of lakes where *A. linnaei* occurred historically or still occurs (summarized in table 2). Information on the pristine state (i.e., the trophic level of a water body without anthropogenic influences) was available for 74 lakes (table 2). Ninety-two percent of these lakes were either oligo- or mesotrophic in their pristine state, indicating typical habitats of *A. linnaei* (with regard to the trophic level) that contrast with van den Hoek’s conclusion, above. For 61 lakes, information on both the pristine state and the current trophic level could be compared. Of these lakes, 92% were originally oligo- or mesotrophic, but more than half of those 61 lakes (57%) are currently eutrophic; therefore, dramatic changes have occurred in the recent past in many habitats of *A. linnaei*. When looking at all lakes with available information on the current trophic state ($n = 80$), the numbers are even more dramatic. Sixty-six percent of the lakes are now eutrophic, or had been in the recent past, and *A. linnaei* still occurs in only 39% of those 80 lakes. These numbers correspond to the general findings presented in figure 2, and strongly suggest that *A. linnaei* occurs mainly in oligomesotrophic habitats, and that eutrophication is correlated with the observed decline of *A. linnaei* populations.

Seemingly contrary to this inferred habitat preference is the finding of populations of *A. linnaei* in 15 eutrophic locations (each addressed below). However, in five of these locations (Takkobu marsh, Pon swamp, Kimoma swamp, Lake Akan, and Boven Wijde; see S3 and S4 at www.nationaalherbarium.nl/supplements/boedeker/BioScience_2010) a strong decline in population size has been observed (Wakana et al. 2001, 2005, 2006, and Boedeker and Immers 2009, respectively), and the future of these populations must be regarded as uncertain. Generally, most eutrophic lakes are turbid and have a poor underwater light climate, but eutrophic clear-water lakes also exist. Shallow lakes, with abundant submerged macrophyte vegetation, may have very clear water with

Table 2. Trophic conditions of lakes where *Aegagropila linnaei* occurs or occurred historically.

Lakes with information available on the pristine state (n = 74)	Percentage
Oligotrophic (including dystrophic)	37
Oligomesotrophic	8
Mesotrophic	47
Slightly eutrophic	7
Eutrophic	1
Lakes with information available on both the pristine and current state (n = 61)^a	
Pristine state oligotrophic (including dystrophic and oligomesotrophic)	39
Pristine state mesotrophic	53
Pristine state eutrophic (including slightly eutrophic)	8
Current state oligotrophic	20
Current state mesotrophic	23
Current eutrophic	57
Lakes with information available on the current state (n = 80)^a	
Current state oligotrophic	16
Current state mesotrophic	18
Current eutrophic	66
<i>Aegagropila linnaei</i> still present	39

a. If a lake had a history of eutrophication but has been restored to a lower nutrient load, it is filed under the most severe stage of eutrophication in its history.

sparse phytoplankton despite relatively high nutrient loadings in lowland areas with soft rock (Phillips et al. 1978). Biological interactions in clear-water lakes differ markedly from “regular” eutrophic lakes (e.g., Jeppesen et al. 1999). The eutrophic clear-water lake Mývatn (Iceland) is well known for its population of *A. linnaei* balls (Einarsson et al. 2004). In Germany, *A. linnaei* has been found in a couple of eutrophic clear-water lakes (Neuklostersee and Teterower See). In Lake Biwa and Lake Kawaguchi (both in Japan), only restricted parts of the lake are eutrophic, while large parts are still mesotrophic (Nagasaka et al. 2002). Two brackish locations of *A. linnaei* are classified as (slightly) eutrophic (Pojo Bay, Finland, and Lake Mälaren, Sweden), as well as one river (River Wear, Scotland). The only “regular” eutrophic lakes where *A. linnaei* was found were Lake Ülemiste (Estonia) and Lake Tiefwareensee (Germany), but the latter had been recently restored to mesotrophic conditions. Even though the evidence is correlative, these numbers strongly suggest that *A. linnaei* occurs mainly in oligomesotrophic habitats but can persist in eutrophic, clear-water lakes, and that indirect effects of eutrophication caused the observed loss of *A. linnaei* populations.

The possible role of lake acidification in the decline of *A. linnaei*

Other factors besides eutrophication may play a role in the decline of *A. linnaei*, such as the widespread and serious problem of lake acidification. Acidification is caused mainly

by the transfer of airborne pollutants by wind from heavily industrialized centers to distant areas where they precipitate as acid rain. Especially in areas where lakes have a low buffering capacity, the effects of acid rain can be drastic, with pH values dropping below 5. Acidification can be a primary and serious problem resulting from airborne dispersal of acidifying pollutants in habitats in remote areas that are not affected by agricultural eutrophication. Many lakes in eastern North America, Great Britain and Ireland, and Scandinavia (except southern Sweden) are characterized by low calcium values and thus a low buffering capacity, which makes them very sensitive to acidification (Jørgensen 1997, Arts 2002). The vegetation of those soft-water lakes is highly endangered, and 90% of the habitats of soft-water vegetation in the Atlantic region have already disappeared (Brouwer et al. 2002). Typical soft-water environments are oligotrophic, with a pH between 5.5 and 7, and have a low buffering capacity (Arts 2002). However, *A. linnaei* is not a typical soft-water organism; instead, the vast majority of habitats are well buffered and characterized by moderate to high calcium levels and an alkaline pH between 7 and 9. No records exist from known acidic habitats. Generally, even less information is available on (historical) acidity than on trophic levels. In the course of this study, only five habitats of *A. linnaei* were found with a pH below 7 (the lowest known pH is 6.3; historical data for Lake Bülte, Germany), and only eight lakes were reported to have low calcium levels. This indicates that *A. linnaei*

typically occurs in habitats that are not strongly affected by acidification. Furthermore, most lakes in Scandinavia have naturally low conductivity, low calcium content, and low pH (Henriksen et al. 1988, Jørgensen 1997), which could possibly explain the absence of *A. linnaei* in Norway and Finland (figure 3b).

Acidification has been shown to be a serious problem in remote mountain lakes that are otherwise unaffected by direct pollution (Curtis et al. 2005). Although several lakes in the Alps region in which *A. linnaei* was historically found were clearly subject to eutrophication (e.g., Ammersee, Lago di Como, Lago Maggiore, Lago di Varese, Starnberger See), it cannot be excluded that acidification might pose an additional threat in more remote mountain areas in noncalcareous parts of the Alps, the Pyrenees, or central Asia. Likewise, acidification might negatively affect *A. linnaei* populations in otherwise unpolluted lakes elsewhere, and thus pose an additional threat of eutrophication in some regions. However, no conclusive data are currently available.

Which effects of eutrophication might cause the observed decline?

Competitive exclusion in the face of increased nutrient loading appears to be the main mechanism responsible for the local decline or even extinction of species in temperate freshwater ecosystems (Barker et al. 2008); that is, species better adapted to high trophic levels outcompete others when nutrient loads increase. It had been proposed that high levels of phosphorus have a toxic effect on charophytes (Forsberg 1965), but this was not confirmed in other studies (Blindow 1992). Even though direct deleterious effects of high nutrient availability are discussed in terrestrial ecosystems, changes in the light climate of aquatic systems seem to be the most important (Smith et al. 1999). The finding of *A. linnaei* in several eutrophic clearwater lakes (see above) could be an indicator that the increase of nutrient levels itself does not negatively affect *A. linnaei*. Shading by increased phytoplankton densities or by opportunistic microalgal epiphytes might be especially significant for *A. linnaei* as a very slow-growing species (Acton 1916, van den Hoek 1963). However, this species survives relatively long and dark winters under the ice in high-latitude lakes such as in Iceland (Jónsson 1992, Einarsson et al. 2004), and healthy balls of *A. linnaei* were found below the shallow photic zone in Lake Teterow, Germany (Pankow and Bolbrinker 1984). This illustrates the capacity of this species to cope with low-light conditions, suggesting that other factors than light limitation might play a more important role in the decline of *A. linnaei* populations.

The attached growth form has been found in several eutrophicated environments such as the Seine in Paris (van den Hoek 1963), Lake Boven Wijde in the Netherlands (Boedeker and Immers 2009), and several Japanese swamp lakes (Wakana et al. 2001, 2005). It is possible for the attached form to survive if it grows on rocks or bivalves close enough to the surface to still receive sufficient light while the water

body becomes more turbid. In Lake Mývatn in Iceland, it has been proposed that variation in the size of submerged *A. linnaei* mats is negatively correlated with (shading) blooms of the cyanobacterium *Anabaena flos-aquae* (Einarsson et al. 2004). Toxins released by blooming cyanobacteria could also be responsible for or contribute to the decline of *A. linnaei* in eutrophicated lakes. Other phenomena correlated with eutrophication that could be linked to the decline of *A. linnaei* are (a) increased feeding of herbivorous fish such as bream, carp, and roach (ten Winkel and Meulemans 1984); (b) local benthic anoxia; (c) higher levels of herbicides and pesticides; (d) greater organic sedimentation; and (e) increased mechanical damage and turbidity caused by recreational boats.

Threat assessment and conservation

Although the exact effects of eutrophication that cause a decline in populations of *A. linnaei* are unclear and require further experimental studies, it seems evident that the decline is correlated with eutrophication. This of course is a general problem that is not easily solved, since the process is closely linked to agriculture, tourism, and human demand and population growth, and is therefore linked to politics and economies. In addition, the populations of *A. linnaei* in several regions and habitats may face specific local problems; for instance, the acid rain degradation of unpolluted, remote habitats. The most famous location of lake balls, Lake Akan, in Japan, is struggling not only with eutrophication and general pollution but also the collection of *Aegagropila* balls as souvenirs. This practice had become so popular that in 1977 the species was classified as critically endangered, and the government asked people to return collected balls to the lake (Soejima et al. 2008, TMLA 2009). Since the populations from the Shatsk lakes in Ukraine seem to be the sole source of the popular *A. linnaei* balls for the entire aquarium trade, these populations must be under considerable, if not extreme, harvesting pressure, especially considering the slow growth rates and longevity of the species (see van den Hoek 1963). In this context it is interesting to note that a Canadian study on nonindigenous plants showed *A. linnaei* to be one of the most common species in the Saint Lawrence Seaway (Cohen et al. 2007), most likely anthropogenically introduced balls originating from amateur aquaria.

Very few molecular studies on *A. linnaei* exist to date. Recently, it has been shown that Lake Akan contains most of the genetic diversity of *A. linnaei* in Japan, as inferred by isozyme studies; therefore Lake Akan is the most important target of conservation efforts (Soejima et al. 2008). Loss of genetic diversity might have occurred already through habitat degradation and extinction of local populations, and further reduction in genetic diversity is to be expected. In many areas, formerly eutrophicated water bodies have been restored to more-or-less natural conditions (van Liere and Gulati 1992, de Bernardo and Giussani 1995). Even after restoration, ecosystems do not necessarily return to their original diversity and community structure (Entwisle

1997). Because of the absence of desiccation-resistant stages in *A. linnaei*, in combination with the species' slow growth and the rarity of spore formation (Leliaert and Boedeker 2007), this species must be regarded as a poor disperser and a poor competitor. Therefore, recolonization of restored habitats might not be very likely. Once a habitat is successfully recolonized by *A. linnaei*, it might take decades before populations are large enough to develop into the typical unattached mats or lake balls. Even if a restored habitat were recolonized, genetic diversity might still be lost. Most extant populations must be regarded as threatened by eutrophication, except in some remote locations. The widespread attached form of *A. linnaei* in the northern Baltic Sea seems to be a stable component of the local algal flora, even though eutrophication has also taken place in this habitat. It is possible that reduced competition in brackish environments plays a role in maintaining *A. linnaei* populations.

One hopes that the popularity of lake balls will help with the conservation of this species and raise awareness of threatened freshwater algae. In addition, we would like to encourage the collection and submission of specimens to accessible collections, not just of *A. linnaei*, but also of species that are less showy and that currently receive less attention.

Acknowledgments

We would like to thank the curators of the herbaria that provided information or access to their collections (S2), as well as Jenny Ask, Leonid V. Averyanov, Thomas Bach, Mandy Bahnwart, Roland Bengtsson (especially for providing figure 1b), Lena Bergström, Hubert Blatterer, Peter Bolbrinker, Ad Bouman, Juliet Brodie, Dwight de Vries, Stefano G. A. Draisma, Árni Einarsson, Tim Entwisle, Peter Feuerpfeil, Olga Gerasymova, Lutz Höhne, Timm Kabus, Maria Kahlert, Kristo K. Kulju, Frederik Leliaert, Inge Lennmark, Eli-Anne Lindstrøm, Liu Guoxiang, Jürgen Mathes, Dieter Mollenhauer, Riggert Munsterhjelm, Alan Pentecost, Bruno Polli, Willem Prud'homme van Reine, Niels Raes, Robert Riegman, Jan Rueness, Clare Scanlan, Volker Schmidt, Hendrik Schreiber, Stephen Skinner, Boris Sviridenko, Maria Thannhauser, Fedor P. Tkachenko, Anna Tolstoy, Bertie J. van Heuven, Kirsten Vendrig, Maria Vorontsova, Isamu Wakana, and Brian A. Whitton. This study was supported in part with funds to CB from the European Union SYNTHESYS program and the Schuure-Beijerinck-Popping Fonds.

References cited

Acton E. 1916. On the structure and origin of "Cladophora balls." *New Phytologist* 15: 1–10.

Arts GHP. 2002. Deterioration of Atlantic soft water macrophyte communities by acidification, eutrophication and alkalisation. *Aquatic Botany* 73: 373–393.

Barker T, Hatton K, O'Connor M, Connor L, Moss B. 2008. Effects of nitrate load on submerged plant biomass and species richness: Results of a mesocosm experiment. *Fundamental and Applied Limnology/Archiv für Hydrobiologie* 173: 89–100.

Bayly IAE, Williams WD. 1973. *Inland Waters and Their Ecology*. Longman.

Bergström L, Bergström U. 1999. Species diversity and distribution of aquatic macrophytes in the Northern Quark, Baltic Sea. *Nordic Journal of Botany* 19: 375–383.

Blindow I. 1992. Decline of charophytes during eutrophication: Comparison with angiosperms. *Freshwater Biology* 28: 9–14.

Boedeker C, Immers I. 2009. No more lake balls (*Aegagropila linnaei* Kützinger, Cladophorophyceae, Chlorophyta) in the Netherlands? *Aquatic Ecology* 43: doi:10.1007/s10452-009-9231-1

Brand F. 1902. Die *Cladophora*-Aegagropilen des Süßwassers. *Hedwigia* 41: 34–71.

Brodie J, John DM, Tittley I, Holmes MJ, Whitton DB. 2008. Important plant areas for algae. (7 January 2010; www.plantlife.org.uk/uk/plantlife-saving-species-publications.html)

Brouwer E, Bobbink R, Roelofs JGM. 2002. Restoration of aquatic macrophyte vegetation in acidified and eutrophied softwater lakes: An overview. *Aquatic Botany* 73: 405–431.

Coesel PFM, Kwakkestein R, Verschoor A. 1978. Oligotrophication and eutrophication tendencies in some Dutch moorland pools, as reflected in their desmid flora. *Hydrobiologia* 61: 21–31.

Cohen J, Mirotchnick N, Leung B. 2007. Thousands introduced annually: The aquarium pathway for non-indigenous plants to the St. Lawrence Seaway. *Frontiers in Ecology and the Environment* 5: 528–532

Curtis CJ, et al. 2005. Acidification in European mountain lake districts: A regional assessment of critical load exceedance. *Aquatic Sciences* 67: 237–251.

de Bernardo R, Giussani G. 1995. Guidelines of Lake Management, vol. 7: Biomaniipulation. Kusatsu-shi. International Lake Environment Committee.

Einarsson Á, Stefánsdóttir G, Jóhannesson H, Ólafsson JS, Gíslason GM, Wakana I, Gudbergsson G, Gardarsson A. 2004. The ecology of Lake Mývatn and the River Laxá: Variation in space and time. *Aquatic Ecology* 38: 317–348.

Duarte CM. 1995. Submerged aquatic vegetation in relation to different nutrient regimes. *Ophelia* 41: 87–112.

Entwisle TJ. 1997. Algae. Pages 34–48 in Scott GAM, Entwisle TJ, May TW, Stevens GN, eds. *A Conservation Overview of Australian Non-marine Lichens, Bryophytes, Algae and Fungi*. Environment Australia.

[EAI] Environment Agency of Japan. 2000. Threatened Wildlife of Japan: Red Data Book, vol. 9: Bryophytes, Algae, Lichens, Fungi. (7 January 2010; www.biodic.go.jp/english/rdb/rdb_e.html)

Forsberg C. 1965. Nutritional studies of *Chara* in axenic cultures. *Physiologia Plantarum* 18: 275–290.

Gardavský A. 1986. *Cladophora moravica* (Dvořák) comb. nova and *C. basiramosa* Schmidle, two interesting riverine species of the green filamentous algae from Czechoslovakia. *Archiv für Hydrobiologie Supplemente* 73: 49–77.

Gärdenfors U. 2005. The 2005 Red List of Swedish Species. Artdatabanken.

Geissler U. 1988. Some changes in the flora and vegetation of algae in freshwater environments. *Helgoländer Meeresuntersuchungen* 42: 637–643.

Getzen MV. 1967. De *Cladophora globosa* in lacubus regionis transpolaris. *Novitates Systematicae Plantarum Non Vascularium* 1: 86–91.

Gíslason GM. 1992. Information Sheet on Ramsar Wetlands. Ramsar Bureau. (11 February 2010; www.wetlands.org/reports/ris/3IS001en_part1.pdf)

Green AJ, Figuerola J, Sánchez MI. 2002. Implications of waterbird ecology for the dispersal of aquatic organisms. *Acta Oecologica* 23: 177–189.

Henriksen A, Lien L, Traaen TS, Sevaldrud IS, Brakke DF. 1988. Lake acidification in Norway: Present and predicted chemical status. *Ambio* 17: 259–266.

Holmes NTH, Whitton BA. 1975. Macrophytes of the River Tweed. *Transactions of the Botanical Society of Edinburgh* 42: 369–381.

———. 1977. Macrophytic vegetation of the River Swale, Yorkshire. *Freshwater Biology* 7: 545–558.

———. 1981. Phytobenthos of the River Tees and its tributaries. *Freshwater Biology* 11: 139–168.

Jeppesen E, Jensen JP, Søndergaard M, Lauridsen T. 1999. Trophic dynamics in turbid and clearwater lakes with special emphasis on the role of zooplankton for water clarity. *Hydrobiologia* 408/409: 217–231.

- John DM. 2002. Order Cladophorales (Siphonocladales). Pages 468–470 in John DM, Whitton BA, Brook AJ, eds. *The Freshwater Algal Flora of the British Isles. An Identification Guide to Freshwater and Terrestrial Algae*. Cambridge University Press.
- Jónsson GS. 1992. Photosynthesis and production of epilithic algal communities in Thingvallavatn. *Oikos* 64: 222–240.
- Jørgensen SE. 1997. Acidification of the Scandinavian lakes. Pages 23–38 in Jørgensen SE, Matsui S, eds. *Guidelines of Lake Management*, vol. 8: *The World's Lakes in Crisis*. International Lake Environment Committee.
- Kann E, Sauer F. 1982. Die "Rotbunte Tiefenbiocönose" Neue Beobachtungen in österreichischen Seen und eine zusammenfassende Darstellung. *Archiv für Hydrobiologie* 95: 181–195.
- Kindle EM. 1915. Limestone solution on the bottom of Lake Ontario. *American Journal of Science* 4: 651–656.
- . 1934. Concerning "lake balls," "*Cladophora* balls" and "coal balls." *American Midland Naturalist* 15: 752–760.
- Kurogi ME. 1980. Lake ball "marimo" in Lake Akan. *Japanese Journal of Phycology* 28: 168–169.
- Leliaert F, Boedeker C. 2007. Cladophorales. Pages 131–183 in Brodie J, Maggs CA, John DM, eds. *Green Seaweeds of Britain and Ireland*. British Phycological Society.
- Lilleleht V. 1998. Red Data Book of Estonia. Threatened Fungi, Plants and Animals. (7 January 2010; www.zbi.ee/punane/english/index.html)
- Ludwig G, Schnittler M. 1996. Rote Liste gefährdeter Pflanzen Deutschlands. Bundesamt für Naturschutz.
- Luther H. 1951. Verbreitung und Ökologie der höheren Wasserpflanzen im Brackwasser der Ekenäs-Gegend in Südfinnland. II. Spezieller Teil. *Acta Botanica Fennica* 50: 1–370.
- [MEA] Millenium Ecosystem Assessment. 2005. *Ecosystems and Human Well-being: Synthesis*. Island Press.
- [MINPRIRODA] Ministry of Natural Resources and Environmental Protection of the Republic of Belarus. 2006. (7 January 2010; www.redbook.minpriroda.by)
- Mollenhauer D. 1998. Gefährdungsursachen für Algen und höhere Wasserpflanzen sowie Möglichkeiten, diese Pflanzen zu erhalten. *Schriftenreihe für Vegetationskunde* 29: 205–217.
- Mollenhauer D, Bengtsson R, Lindström E-A. 1999. Macroscopic cyanobacteria of the genus *Nostoc*: A neglected and endangered constituent of European inland aquatic biodiversity. *European Journal of Phycology* 34: 349–360.
- Moriwake N. 2005. Information Sheet on Ramsar Wetlands. Ramsar Bureau. (12 January 2010; www.wetlands.org/reports/ris/2JP014_RISen05.pdf)
- Nagasaka M, Yoshizawa K, Ariizumi K, Hirabayashi K. 2002. Temporal changes and vertical distribution of macrophytes in Lake Kawaguchi. *Limnology* 3: 107–114.
- Nakazawa S. 1974. The time and the cause of extermination of lake balls from Lake Zeller. *Bulletin of the Japanese Society of Phycology* 12: 101–103.
- Noskov GA. 2000. Red Data Book of Nature of the Leningrad Region, vol. 2. St. Petersburg.
- Palik P. 1963. Studien über *Aegagropila sauteri* (Neetz) Kützing. *Annales Universitatis Scientiarum Budapestinensis Sectio Biologica* 5: 179–192.
- Pankow H. 1965. *Aegagropila sauteri* in Mecklenburg (Norddeutschland). *Nova Hedwigia* 9: 177–184.
- . 1985. Verschollene, gefährdete und interessante Großalgen im nördlichen Gebiet der DDR. *Botanische Rundbriefe des Bezirks Neubrandenburg* 16: 65–72.
- Pankow H, Bolbrinker B. 1984. Über die Verbreitung und Soziologie von *Cladophora aegagropila* (L.) Rbh [= *Aegagropila sauteri* (Nees ex Kütz.) Kütz.] in den Nordbezirken der DDR. *Gleditschia* 12: 279–283.
- Phillips GL, Eminson D, Moss B. 1978. A mechanism to account for macrophyte decline in progressively eutrophicated freshwaters. *Aquatic Botany* 4: 103–126.
- Revinga C, Campbell I, Abell R, de Villiers P, Bryer M. 2005. Prospects for monitoring freshwater ecosystems toward the 2010 targets. *Philosophical Transactions of the Royal Society B* 360: 397–413.
- Schlichting HE. 1960. The role of waterfowl in the dispersal of algae. *Transactions of the American Microscopical Society* 79: 160–166.
- Smith VH, Tilman GD, Nekola JC. 1999. Eutrophication: Impacts of excess nutrient inputs on freshwater, marine, and terrestrial ecosystems. *Environmental Pollution* 100: 179–196.
- Soejima A, Yamazaki N, Nishino T, Wakana I. 2008. Genetic variation and structure of the endangered freshwater benthic alga *Marimo*, *Aegagropila linnaei* (Ulvophyceae) in Japanese lakes. *Aquatic Ecology* 43: 359–370.
- Taranu ZE, Gregory-Eaves I. 2008. Quantifying relationships among phosphorus, agriculture, and lake depth at an inter-regional scale. *Ecosystems* 11: 715–725.
- ten Winkel EH, Meulemans JT. 1984. Effects of fish upon submerged vegetation. *Hydrobiological Bulletin* 18: 157–158.
- [TMLA] The Marimo of Lake Akan. 2009. Marimo are returned to Lake Akan! (7 January 2010; www.marimo-web.org/en/c-history/c2-marimo_return.html)
- van den Hoek C. 1963. Revision of the European Species of *Cladophora*. E. J. Brill.
- van Liere L, Gulati RD. 1992. Restoration and Recovery of Shallow Eutrophic Lake Ecosystems in the Netherlands. Kluwer Academic Publishers.
- Waern M. 1952. Rocky-shore algae in the Öregrund Archipelago. *Acta phytogeographica Suecica* 30: 1–298.
- Wakana I, Arai S, Sano O, Honoko H. 2001. Current situation and living environment of freshwater alga "Tateyama-marimo" (*Aegagropila* sp. nov.) in the Saryfutsu-Hamatonbetsu lake group, Hokkaido, Japan. *Journal of Plant Research* 114 (suppl.): 59.
- Wakana I, Kawachi M, Hamada N, Taoda H. 2005. Current situation and living environment of an endangered algal species marimo (*Aegagropila linnaei*) in Takkobu marsh, Kushiro-Shitsugen wetland. *Journal of Plant Research* 118 (suppl.): 64.
- Wakana I, Satoh H, Suzuki Y. 2006. The past and the present of vegetation and environment in a disappeared habitat of the freshwater alga marimo (*Aegagropila linnaei*) in Lake Akan, Japan. *Journal of Plant Research* 114 (suppl.): 63.
- Wesenberg-Lund C. 1903. Sur les *Aegagropila sauteri* du Lac de Sorö. *Académie Royale des Sciences et des Lettres de Danemark* 2: 167–203.
- Whitton BA, Boulton PNG, Clegg EM, Gemmill JJ, Graham GG, Gustar R, Moorhouse TP. 1998. Long-term changes in macrophytes of British rivers, 1: River wear. *Science of the Total Environment* 210/211: 411–426.

Christian Boedeker (boedeker@nhn.leidenuniv.nl), Anne Immers, and Erik Smets are with the National Herbarium of The Netherlands, Leiden University Branch, in Leiden, The Netherlands. Anja Eggert works in Physical Oceanography and Instrumentation, at the Leibniz Institute for Baltic Sea Research Warnemünde, in Rostock, Germany.